

Predation on the tiger rat snake *Spilotes pullatus* (Serpentes: Colubridae) by the southern black-eared opossum *Didelphis aurita* (Mammalia: Didelphimorphia), with a review of known predators

Anderson Lozorio Canal Filho^{1*}, Isabella Marçal Silva², Beatriz Medeiros Brandão³, Iasmin Macedo⁴, Thiago Silva-Soares⁵

1 Programa de Pós Graduação em Ciências de Floretas Tropicais, Instituto Nacional de Pesquisas da Amazônia, 69060-001 Manaus, AM, Brazil

2 Universidade Federal do Espírito Santo, 29500-000 Alegre, ES, Brazil

3 Rua Francisco Xavier 144, casa 4, Morro das Pedras, 88066-027 Florianópolis, SC, Brazil

4 Instituto Últimos Refúgios, Rua Amarílio Lunz 16, República, 29070-030 Vitória, ES, Brazil

5 Herpeto Capixaba project, Instituto Biodiversidade Neotropical, Rua Sanhaço 562, Nova Guarapari, 29206-400 Guarapari, ES, Brazil

*Corresponding author: andersonlozorio@gmail.com

DOI: [10.5281/zenodo.10204741](https://doi.org/10.5281/zenodo.10204741)

The snake genus *Spilotes* Wagler, 1830 (Colubridae) is currently composed of two species: *Spilotes pullatus* (Linnaeus, 1758) and *Spilotes sulphureus* (Wagler, 1824) (Jadin, 2013). *Spilotes pullatus* is popularly known in Brazil as “caninana”, which in Tupi language means “that has small head” (Amaral, 1929). This vernacular name refers to the diminutive head size in relation to the large sizes that adult individuals can

attain – up to 270 mm total length – as one of the largest colubrids in South America (Savage, 2002; Marques et al., 2014). One of the main characteristics of *S. pullatus* is its fascinating ability to expand the neck region, due to the elasticity and great extension capacity of a membrane that connects the ends of tracheal rings, which allows the species to appear larger in a threatening situation (Amaral, 1929).

This species, listed as Least Concern in the IUCN Red List (Arzamendia et al., 2019), is widely distributed in Central and South America, from southern Mexico to northern Argentina (Savage, 2002). It is widespread in Brazil, present in all biomes (Vanzolini et al., 1980; Mendonça et al., 2011; Nogueira et al., 2019). It is an arboreal and diurnal species, non-venomous and with a generalist diet, swallowing prey alive or killing them by constriction (Martins et al., 2008; Marques et al., 2014; Castro & Silva-Soares, 2016).

The southern black-eared opossum *Didelphis aurita* Wied, 1826 (Didelphimorphia, Didelphidae) is one of the most common marsupial species in the eastern Neotropical region, occurring from northeastern Brazil to northeastern Argentina, inhabiting mainly primary and secondary forests at lower elevations (Astúa, 2015). It is a scansorial, solitary and nocturnal marsupial inhabiting a wide range of habitats in the Atlantic Forest, including urban and rural areas (Cerqueira et al., 1990; Vieira & Monteiro-Filho, 2003). The characteristics associated with its generalist food habits (opportunistic/omnivorous) make this species largely abundant, with high population densities, especially in areas close to human habitations, hence, a sylvatic-synanthropic species (Olifiers et al., 2005; Junior & Leite 2007).

On 20 February 2022 at 1912h, an adult *Spilotes pullatus* was filmed in an attempted predation by an adult *D. aurita*. The interaction was recorded at Ilhabela, an archipelago off the northern coast of the state of São Paulo, southeastern Brazil (23°49'14.43" S 45°21'17.58" W, datum WGS84; 26 m elev. a.s.l.), adjacent to an urban area near Ilhabela State Park, a 27,025 hectare "archipelago-park" covered by Atlantic Ombrophilous Dense Forest.

The interaction occurred as follows: BMD heard a loud crawling sound in leaf litter. The *S. pullatus* emerged from the forest, pursued five seconds later by the *Didelphis aurita*. The observer began videotaping the scene (Figure 1). The opossum rapidly attacked the snake, which tried to strike and bite the marsupial once or twice. The snake also attempted to coil its body around the opossum, but the marsupial shook off the snake. The opossum did not immobilize the snake on its first attack, but bit the snake behind the head, immobilizing it and carrying it, still alive, in its mouth back towards the forest. The video is 21 seconds in length, but the total interaction, from the first noise to the animals' disappearance, lasted about 35 seconds. The video may be accessed at herpetocapixaba.com.br/herpetovideos.

Didelphis aurita locates food items mainly through auditory and olfactory

stimuli, usually grasping the food with its mouth directly from the ground (Hunsaker, 1977; Streilein, 1982). The opossum and the caninana are sympatric throughout their distributions and most species of *Didelphis* are known to prey upon snakes (Cordero & Nicolas, 1986; Santori et al., 1995; Astúa, 2015). Since *S. pullatus* is diurnal and *D. aurita* is nocturnal, we believe that the opossum probably encountered the snake while it was resting in a nocturnal shelter. Since snake predators typically consume prey that are smaller than the predator (Schalk & Cove, 2018), it is remarkable that the opossum attacked and subdued the snake, which measured approximately 200 cm. Despite these records, *D. aurita* is apparently not a specialized snake predator, and the action of attacking a snake by a strong bite at the base of the head followed by shaking it from side to side, is apparently a conservative behavior in mammals (Oliveira & Santori, 1999).

To search for records of *Didelphis aurita* preying on *S. pullatus*, we performed a literature review through the Web of Science, Scopus and Google Scholar online platforms. We incorporated search terms to cover literature published in English, Portuguese, and Spanish, followed OR by “*Spilotes pullatus*” or “*Didelphis aurita*”. In addition, a search was made for all available issues of the journal Herpetological Review (1967-2022). Finally, we searched

for photographic snake feeding records on the Wiki Aves platform on 26 March 2023 with the help of the online material available by Souza et al., (2022) in attempt to add more records. Although documented predation records for *S. pullatus* are scarce, the species has been reported as prey of reptiles, birds, and mammals (Tab. 1).

Most studies on the interaction of *Didelphis* opossums and snakes have focused on venomous snakes, since it is documented that at least four species of *Didelphis* are tolerant to snake venom (Vellard, 1945; Perales et al., 1986; Oliveira & Santori, 1999; Astúa, 2015). We found records of *S. pullatus* being preyed by raptors such as the Great Black Hawk – *Urubitinga urubitinga* (Gmelin, 1788), the Roadside Hawk – *Rupornis magnirostris* (Gmelin, 1788) and by the specialized snake-eater, the Laughing-Falcon – *Herpetotheres cachinnans* (Linnaeus, 1758) (Gerhardt et al., 1993; Costa et al., 2014; Oakley et al., 2021). Furthermore, a predation attempt by the Central American Indigo Snake *Drymarchon melanurus* (Duméril, Bibron & Duméril, 1854) was also reported (Oakley & Theodorou, 2020) and previously, Hernández-Ríos et al., (2013) observed a *D. melanurus* in the process of consuming a *S. pullatus* in Mexico; however, because the initial capture of the prey was not observed, it is unknown if the *S. pullatus* was preyed or scavenged. More recent-

ly, Peláez-Cruz et al. (2022) reported a potential predation event on *S. pullatus* by the gray fox *Urocyon cinereoargenteus* (Schreber, 1775) recorded in northern Yucatán, Mexico, although there is no evidence that the fox consumed the snake.

After searching these sources, we conclude that predators of *S. pullatus* are poorly documented. The predator-prey interaction between *D. aurita* and *S. pullatus* has not been previously reported, so this is the first record of such predation, with the description of their struggle. Our record adds one more potential prey of *Didelphis aurita*, as well as one more predator of *Spilotes pullatus*, showing the likely opportunistic behavior of the marsupial. This new record broadens our knowledge about the natural history of the species, and we suggest considering the tiger rat snake as a component of the opossum diet.

ACKNOWLEDGMENTS

We thank Marsupial Project, Instituto Últimos Refúgios, for all partnership along the past years. This work is part of 'Herpeto Capixaba: for the knowledge and conservation of amphibians and reptiles of Brazil' and was partially financed in part by Fundação de Amparo à Pesquisa e Inovação do Espírito Santo (EDITAL FAPES Nº 03/2021 - UNIVERSAL #437/2021). We also thank Henrique C. Costa for the suggestions and comments on the manuscript.

REFERENCES

Amaral A.D. 1929. Estudo sobre ophidios neotrópicos XIX. Revisão do gênero *Spilotes* Wagler, 1830. *Memórias do Instituto Butantan* 4:275–298.

Arzamendia V., Caicedo J.R., Fitzgerald, L., Giraud, A., Gutiérrez-Cárdenas, P., Kacolis... Murphy, J. 2019. *Spilotes pullatus*. The IUCN Red List of Threatened Species 2019. Accessed on 08 February 2023.

Astúa D. 2015. Family Didelphidae (Opossums). Pp. 70–186, in Wilson D.E., Mittermeier R.A. (Eds.), *Handbook of the Mammals of the World* . v.5. Monotremes and Marsupials. Lynx Edicions, Barcelona.

Boos H.E. 2001. The snakes of Trinidad and Tobago. Texas A&M University Press.

Castro T.M., Silva-Soares T. 2016. Répteis da restinga do Parque Estadual Paulo César Vinha: Guarapari, Espírito Santo, Sudeste do Brasil. Centro Universitário São Camilo, Cachoeiro do Itapemirim.

Cerqueira R., Fernandez F.A.S., Quintela M.F.S. 1990. Mamíferos da restinga de Barra de Maricá, Rio de Janeiro. *Papéis Avulsos de Zoologia* 37:141–157.

- Champagne P., Singer D. 2021. *Clelia clelia* (Black-headed mussurana). Diet & Prey. *Herpetological Review* 52:865.
- Cordero G.A.R., Nicolas R.A.B. 1986. Feeding habits of the opossum (*Didelphis marsupialis*) in northern Venezuela. *Fieldiana Zoology* 39:125–131.
- Costa H.C., Lopes L.E., Freitas M.B., Zorzin G. 2014. The reptile hunter's menu: A review of the prey species of Laughing Falcons, *Herpetotheres cachinnans* (Aves: Falconiformes). *North-Western Journal of Zoology* 10:445–453.
- Gans, C. 1986. Functional morphology of predator-prey relationships. Pp. 6–23, in Feder, M.E., Lauder G.V, (Eds.), *Predator-prey relationships*. University of Chicago, Chicago.
- Gerhardt R.P., Harris P.M., Marroquin M.A.V. 1993. Food habits of nesting great black hawks in Tikal National Park, Guatemala. *Biotropica* 25:349–352. doi: 10.2307/2388794.
- Hernández-Ríos A., García-Padilla E., Villagas-Nieto A. 2013. *Drymarchon melanurus* (Central American Indigo Snake). Diet. *Herpetological Review* 44:690.
- Hunsaker I.I.D. 1977. Ecology of New World marsupials. Pp. 95–156, in: Hunsaker I.I.D. (Ed.), *The biology of marsupials*. Academic Press, New York.
- Jadin R.C. 2013. Untangling a phylogenetic knot of serpents: Integrative systematics on Neotropical snakes. Ph. D. dissertation, University of Colorado.
- Junior V.C., Leite Y.L.R. 2007. Uso de habitats por pequenos mamíferos no Parque Estadual da Fonte Grande, Vitória, Espírito Santo, Brasil. *Boletim do Museu de Biologia Mello Leitão* 21:57–77.
- Marques O.A., Muniz-da-Silva D.F., Barbo F.E., Cardoso S.R.T., Maia D.C., Almeida-Santos S.M. 2014. Ecology of the Colubrid snake *Spilotes pullatus* from the Atlantic Forest of Southeastern Brazil. *Herpetologica* 70:407–416. doi: 10.1655/HERPETOLOGICA-D-14-00012.
- Marques O.A.V., Abe A.S., Martins, M. 1998. Estudo diagnóstico da diversidade de répteis do Estado de São Paulo. Biodiversidade do Estado de São Paulo, Brasil: síntese do conhecimento ao final do século XX. Editora Fapesp, São Paulo.
- Martins M., Marques O.A., Sazima, I. 2008. How to be arboreal and diurnal and still stay alive: microhabitat use, time of activity, and defense in Neotropical forest snakes. *South American Journal of Herpetology* 3:58–67.

- Mendonça P.P., Cobra P., Bernardo L.R., Silva-Soares, T. 2011. Predation of the snake *Spilotes pullatus* (Squamata: Serpentes) upon the rodent *Proechimys gardneri* (Rodentia: Echimyidae) in the Amazonian basin, northwestern Brazil. *Herpetology Notes* 4:425–427.
- Nogueira C.C., Argôlo A.J.S., Arzamendia V., Azevedo J.A., Bérnils R.S., Bolochio, B.E., ... Martins M. 2019. Atlas of Brazilian snakes: verified point-locality maps to mitigate the wallacean shortfall in a megadiverse snake fauna. *South American Journal of Herpetology* 14:1–274. doi: 10.2994/SA-JH-D-19-00120.1.
- Novy S.A., Van Putte R.D. 2016. Behavioral Notes and Nesting of the Black Solitary Eagle (*Buteogallus solitarius*) in Belize. *Transactions of the Illinois State Academy of Science* 109:29–33.
- Oakley J., Theodorou A. 2020. A failed predation attempt by a Central American Indigo Snake (*Drymarchon melanurus*) on a Tiger Ratsnake (*Spilotes pullatus*) in Campeche, Mexico. *Reptiles & Amphibians* 27:494–495. doi: 10.17161/randa.v27i3.14895.
- Oakley J., Theodorou A., Adojaan A., Vallocchia S. 2021. *Spilotes pullatus* (Tiger Ratsnake). Predation. *Herpetological Review* 51:360.
- Olifiers N., Gentile R., Fiszon J.T. 2005. Relation between small-mammal species composition and anthropic variables in the Brazilian Atlantic Forest. *Brazilian Journal of Biology* 65:495–501. doi:10.1590/S1519-69842005000300015.
- Oliveira M.E., Santori R.T. 1999. Predatory behavior of the opossum *Didelphis albiventris* on the pitviper *Bothrops jararaca*. *Studies on Neotropical Fauna and Environment* 34:72–75. doi: 10.1076/snfe.34.72.2105.
- Perales J., Muños R., Moussatche, H. 1986. Isolation and partial characterization of a protein fraction from the opossum (*Didelphis marsupialis*) serum, with protecting property against the *Bothrops jararaca* venom. *Anais da Academia Brasileira Ciências* 58:155–162.
- Pontes J.A.L., Figueiredo J.P., Pontes R.C., Rocha C.F.D. 2008. Snakes from the Atlantic Rainforest area of Serra do Mendanha, in Rio de Janeiro state, southeastern Brazil: a first approximation to the taxocenosis composition. *Brazilian Journal of Biology* 68:601–608. doi: 10.1590/S1519-69842008000300018.
- Santori R.T., Moraes D.A., Cerqueira R. 1995. Diet composition of *Metachirus nudicaudatus* and *Didelphis aurita* (Marsupialia, Didelphoidea) in South-

eastern Brazil. *Mammalia* 59: 511–516. doi: [10.1515/mamm.1995.59.4.511](https://doi.org/10.1515/mamm.1995.59.4.511).

Savage J.M. 2002. The amphibians and reptiles of Costa Rica: a herpetofauna between two continents, between two seas. Academic Press, University of Chicago.

Sazima I., Haddad C.F.B. 1992. Répteis da Serra do Japi: notas sobre história natural. Pp. 28–49, in Morellato, L.P. (Ed.), História natural da Serra do Japi: ecologia e preservação de uma área florestal no sudeste do Brasil.

Schalk C.M., Cove M.V. 2018. Squamates as prey: Predator diversity patterns and predator-prey size relationships. *Food Webs* 17:e00103. doi: doi.org/10.1016/j.fooweb.2018.e00103.

Silva J.S., El-Deir A.C., Moura G.J., Alves R.R., Albuquerque U.P. 2014. Traditional ecological knowledge about dietary and reproductive characteristics of *Tupinambis merianae* and *Hoplias malabaricus* in Semiarid Northeastern Brazil. *Human Ecology* 42:901–911. doi: doi.org/10.1007/s10745-014-9698-9.

Souza E., Lima-Santos J., Entiuspe-Neto O. M., Santos M. M., Moura P.R., Hingst-Zaher E. 2022. Ophiophagy in Brazilian birds: a contribution from a collaborative platform of citizen

science. *Ornithology Research* 30:15–24. doi:[10.1007/s43388-022-00082-5](https://doi.org/10.1007/s43388-022-00082-5).

Streilein K.E. 1982: Behavior, ecology, and distribution of South American marsupials. *Special Publication of the Pymatuning Laboratory of Ecology* 6: 231–250.

Uetz, P., Freed, P, Aguilar, R. & Hosek, J. (Eds.). 2022. The Reptile Database: <http://www.reptile-database.org>. Accessed on 02 November 2022.

Vanzolini P.E., Ramos-Costa A.M.M., Vitt, L.J. 1980. Répteis das caatingas. Academia Brasileira de Ciências, Rio de Janeiro.

Velame R. 2013. [WA1122818, *Carcara plancus* (Miller, 1777)]. Wiki Aves - The Encyclopedia of Birds of Brazil. Accessible at <http://www.wikia.ves.com/1122818>. Accessed on 26 March 2023.

Vellard, J. 1945. Resistencia de los “*Didelphis*” (zarigueya) a los venenos ofídicos; nota prévia. *Revista Brasileira de Biologia* 5:463–467.

Vieira E.M., Monteiro-Filho E. L. 2003. Vertical stratification of small mammals in the Atlantic rain forest of south-eastern Brazil. *Journal of Tropical Ecology* 19:501–507. doi: [10.1017/S0266467403003559](https://doi.org/10.1017/S0266467403003559).

Editor: Henrique C. Costa

Table 1. Predators previously reported for the Tiger Rat Snake *Spilotes pullatus*.

(*) Photo Web Link. Velame (2013). Observation code:WA1122818.

PREDATOR	LOCALITY	REFERENCES
SQUAMATA		
LIZARDS		
Teiidae		
<i>Salvator merianae</i> (Duméril & Bibron, 1839)	Ceara, Brazil	Silva et al., (2014)
SERPENTES		
Colubridae		
<i>Drymarchon melanurus</i> (Duméril, Bibron & Duméril, 1854)	Tamaulipas, Mexico	Hernández-Ríos et al., (2013)
	Campeche, Mexico	Oakley & Theodorou (2020)
Dipsadidae		
<i>Clelia clelia</i> (Daudin, 1803)	Tambopata, Peru	Champagne (2021)
AVES		
Accipitriformes		
Accipitridae		
<i>Rupornis magnirostris</i> (Gmelin, 1788)	Campeche, Mexico	Oakley et al., (2021)
<i>Urubitinga solitaria</i> (Vieillot, 1817)	Cayo District, Belize	Novy & Van Putte (2016)
<i>Urubitinga urubitinga</i> (Gmelin, 1788)	Flores, Guatemala	Gerhardt et al., (1993)
Falconiformes		
Falconidae		
<i>Caracara plancus</i> (Miller, 1777)	Bahia, Brazil	https://www.wikiaves.com.br/1122818*
<i>Herpetotheres cachinnans</i> (Linnaeus, 1758)	Costa Rica	Costa et al., (2014)
MAMMALIA		
Carnivora		
Canidae		
<i>Urocyon cinereoargenteus</i> (Schreber, 1775)	Yucatán, Mexico	Peláez-Cruz et al., (2022)
Didelphimorphia		
Didelphidae		
<i>Didelphis aurita</i> (Wied, 1826)	São Paulo, Brazil	This study



Figure 1. Southern black-eared opossum (*Didelphis aurita*) preying on a tiger rat snake (*Spilotes pullatus*), in Ilhabela, state of São Paulo, Brazil: (A-B) Opossum attempts to kill the snake; (C) snake tries to strike and coil its body around the marsupial; (D) the snake is finally immobilized with a bite behind the head. Photos: BMD.