

An unexpected encounter: first record of predation on *Hemidactylus mabouia* (Squamata: Gekkonidae) by a juvenile *Mastigodryas boddaerti* (Squamata: Colubridae)

David Soares^{1,2*}, Tássio Alves Coêlho³, Darlison Chagas-de-Souza³, Antônio Willian L. da Trindade², Lívia Carla Chagas⁴, Síría Ribeiro^{1,2}, Alfredo P. Santos-Jr.²

1 Programa de Educação Tutorial (PET) Conexões de Saberes de Estudos Interdisciplinares: Comunidades de Campo, Universidade Federal do Oeste do Pará, Rua Vera Paz, Bloco Multidisciplinar Tapajós II, S/N Salé, Santarém, Pará, 68040-255, Brazil.

2 Laboratório de Ecologia e Comportamento Animal (LECAN), Universidade Federal do Oeste do Pará Rua Vera Paz S/N Salé, Santarém, Pará, 68040-255, Brazil.

3 Laboratório de Estudos Morfofisiológicos e Parasitários, Departamento de Ciências Biológicas e da Saúde, Universidade Federal do Amapá, Rod. Juscelino Kubitschek, s/n, Jardim Marco Zero, Macapá, Amapá 68903-419, Brazil.

4 Laboratório de Fisiologia e Toxinas Animais – Fístox, Universidade Federal do Oeste do Pará, Avenida Marechal Rondon s/n, Caranazal, Santarém, Pará 68040-070, Brazil.

* Corresponding author. E-mail: davidmarcelo200151@gmail.com

The Amazon region is one of the most biodiverse forests in the world, hosting a wide variety of snake species (Vitt, 1987; Guedes et al., 2023). Despite this high diversity, little is known about the ecological aspects of this fauna, such as feeding biology (Schmidt & Inger, 1951; Hoge & Nina, 1969; Vitt, 1987; Martins, 1993; Caldwell, 1996; Martins & Oliveira, 1998). Amphibians and lizards are common dietary items for many snakes, although they may also

feed on small invertebrates, fish, birds, small mammals, and even other snakes (Duellman, 1978; Martins & Oliveira, 1998). Knowledge of the trophic interactions of snakes would improve the understanding of their natural history and their role in the niche they occupy (Cadle & Greene, 1993; Alencar et al., 2013; Pinto-Coelho et al., 2021).

Mastigodryas boddaerti (Sentzen, 1796) is a snake widely distributed in the Amazon biome, with records in oth-

er biomes (Cerrado, Caatinga and Pantanal) (Nogueira et al., 2019). It occurs in anthropogenic and peri-anthropogenic environments, as well as in primary and secondary forests (Fraga et al., 2013; Guedes et al., 2014). This species is considered a generalist in both habitats and diet, which mainly includes amphibians, lizards, amphisbaenians, birds, and small mammals (Siqueira et al., 2012; Eversole et al., 2023). However, there is little data on its diet during ontogenetic stages (Eversole et al., 2023). This is the first record of the lizard *Hemidactylus mabouia* (Moreau De Jonnès, 1818) as prey of a juvenile *M. boddaerti*.

A juvenile *M. boddaerti* (snout-vent length [SVL] 335 mm; tail length [TL] 124.72 mm) was found dead on a walking trail during a recreational activity in April 2014 in a forest fragment at the Vila Americana plateau site, municipality of Belterra (2°38'19.00"S, 54°56'39.05"W; elevation 166 m, SIRGAS2000), state of Pará, Brazil. The specimen was sent to the Laboratório de Ecologia e Comportamento Animal (LECAN) at the Universidade Federal do Oeste do Pará (UFOPA) and deposited under the accession number UFOPA-H 3352. Examination of the specimen determined that death was caused by ingesting an individual of the wall lizard, *H. mabouia* (SVL 52.07 mm; TL 28.98 mm). The lizard was likely too large for the predator, as its SVL was

approximately 25% of the snake's SVL, causing the snake's stomach to rupture, exposing the lizard (Fig. 1A-C) and subsequently leading to the snake's death.

Hemidactylus mabouia is an exotic lizard from the African continent that was introduced to Brazil on slave ships (Vanzolini, 1978; Carranza & Arnold, 2006; Anjos & Rocha 2008). Today, it is widely distributed in all South American countries, and occurs in all Brazilian biomes (Ribeiro-Júnior, 2015;). The species is nocturnal and feeds on invertebrates. It is associated with anthropogenic, peri-anthropogenic and natural environments, often found on walls in anthropogenic areas and on tree trunks in forest environments (Avila-Pires, 1995; Vitt et al., 2008; Bonfiglio et al., 2006).

Snakes often feed on larger prey, for the cost-benefit of consuming larger specimens, which provides a greater food source (Shine, 1991). The genus *Hemidactylus* was previously recorded as an item in the diet of *M. boddaerti* by Siqueira et al. (2012) but those authors were unable to identify the prey at the species level, since two species occur in the Amazon region: *H. mabouia* and *Hemidactylus palaichthus* Kluge, 1969 (Avila-Pires, 1995).

Occasional encounters such as this can provide important data on the ecology of various vertebrates as well as

predation processes. Also, biological collections are important sources of information on the ecological aspects of different groups of animals and are extremely important for conducting research and obtaining data.

ACKNOWLEDGMENTS

We thank Coordenação de Aperfeiçoamento de Pessoal de Nível Superior for scholarships granted to TAC and DCS (Process 88887.598663/2021-00 and 88887.636892/2021-00, respectively). We would like to thank the Fundação Amazônia de Amparo a Estudos e Pesquisa do Pará (FAPESPA) for the scholarship granted to AWLT. We thank Fundo Nacional de Desenvolvimento e Educação for promoting the scholarships awarded from the Programa de Educação Tutorial-PET to DS.

REFERENCES

- Alencar L.R.V., Gaiarsa M.P., Martins M. 2013. The Evolution of Diet and Microhabitat use in Pseudoboine Snakes. *South American Journal of Herpetology* 8:60–66.
- Anjos L.A., Rocha C.F.D. 2008. The *Hemidactylus mabouia* Moreau de Jonnes, 1818 (Gekkonidae) lizard: an invasive alien species broadly distributed in Brazil. *Natureza & Conservação* 6:196–207.
- Avila-Pires T.C.S. 1995. Lizards of Brazilian Amazonia (Reptilia: Squamata). *Zoologische Verhandelingen* 299:1–706.
- Beebe W. 1946. Field notes on the snakes of Kartabo, British Guiana, and Caripito, Venezuela. *Zoologica* 4:1–52.
- Bernarde P.S., Abe A.S. 2010. Hábitos alimentares de serpentes em Espigão do Oeste, Rondônia, Brasil. *Biota Neotropica* 10:167–173. Doi: 10.1590/S1676-06032010000100017.
- Bonfiglio F., Balestrin, R.L., Cappellari L.H. 2006. Diet of *Hemidactylus mabouia* (Sauria, Gekkonidae) in urban area of southern Brazil. *Biociências* 14:107–111.
- Cadle J.E., Greene H.W. 1993. Phylogenetic patterns, biogeography, and ecological structure of Neotropical snake assemblages. Pp. 281–293, in Ricklefs R.E., Schluter D. (Eds.), *Species diversity in ecological communities. Historical and geographical perspectives*. University of Chicago Press, Chicago.
- Caldwell J.P. 1996. Diversity of Amazonian anurans: The role of systematics and phylogeny in identifying macroecological and evolutionary patterns. Pp. 73–88, in Gibson A.C. (Ed.). *Neotropical Biodiversity and Conservation* Mildred E. Mathias Botanical Garden. Los Angeles.

- Carranza S., Arnold E.N. 2006. Systematics, biogeography, and evolution of *Hemidactylus geckos* (Reptilia: Gekkonidae) elucidated using mitochondrial DNA sequences. *Molecular Phylogenetics and Evolution* 38:531–545.
- Duellman W.E. 1978. The biology of an equatorial herpetofauna in Amazonian Ecuador. *Miscellaneous Publications - Museum of Natural History, University Kansas* 65:1-352.
- Eversole C.B., Powell R.L., Rivas L.R. 2023. *Mastigodryas boddaerti* (Boddaert's Tropical Racer). Diet. *Herpetological Review* 54:489.
- Forti L., Kawashita-Ribeiro R. 2008. *Mastigodryas boddaerti* (Boddaert's tropical Racer): Predation. *Herpetological Bulletin* 106:38–40.
- Fraga R., Lima A.P., Prudente A.L.C., Magnusson W.E. 2013. Guide to the snakes of the Manaus region - Central Amazonia. Editora INPA, Manaus.
- Guedes T.B., Entiauspe-Neto O.M., Costa H. C. 2023. Lista de répteis do Brasil: Atualização de 2022. *Herpetologia Brasileira* 12:56–161. Doi: 10.5281/ZENODO.7829013.
- Guedes T.B., Nogueira C., Marques O.A.V. 2014. Diversity, natural history, and geographic distribution of snakes in the Caatinga, Northeastern Brazil. *Zootaxa* 3863:1–93.
- Hoge A.R., Nina A.C.M. 1969. Serpentes coletadas pelo Instituto Nacional de Pesquisas da Amazonia. *Memórias do Instituto Butantan* 30:71–96.
- Martins M. 1993. Why do snakes sleep on the vegetation in Central Amazonia? *Herpetological Review* 24:83-84.
- Martins M., Oliveira M.E. 1998. Natural history of snakes in forests of the Manaus region, Central Amazonia, Brazil. *Herpetological Natural History* 6:78–150.
- Nogueira C.C., Argôlo A.J.S., Arzamendia V., Azevedo J.A., Barbo F.E., Bérnils R.S., Martins M.C.M. 2019. Atlas of Brazilian snakes: verified point-locality maps to mitigate the Wallacean shortfall in a megadiverse snake fauna. *South American Journal of Herpetology* 14:1–274.
- Pinto-Coelho D., Martins M., Guimarães-Junior P.R. 2021. Network analyses reveal the role of large snakes in connecting feeding guilds in a species-rich Amazonian snake community. *Ecology and Evolution* 11:6558–6568.
- Ribeiro-Junior M.A. 2015. Catalogue of distribution of lizards (Reptilia: Squamata) from the Brazilian Amazonia. II. Gekkonidae, Phyllodactylidae, Sphaerodactylidae. *Zootaxa* 3981:1–55.

Schmidt K.P., Inger R.F. 1951. Amphibians and reptiles of the Hopkins-Brandner expedition to Brazil. *Fieldiana Zoology* 31:439–465.

Shine, R. 1991. Why do larger snakes eat larger prey items? *Functional Ecology* 5: 493–502.

Siqueira D.M., Nascimento L.P., Santos-Costa M.C. 2012. Feeding biology of boddaert's tropical racer, *Mastigodryas boddaerti* (Serpentes, Colubridae) from the Brazilian Amazon. *South American Journal of Herpetology* 7:226–232.

Vanzolini P.E. 1978. An annotated bibliography of the land and fresh-water reptiles of South America. *Museu de Zoologia* 2:1758-1975. doi: 10.5962/bhl.title.100694.

Vitt L.J. 1987. Communities. Pp. 335–365, in Eigel R.A.S., Colins J.T., NOVAK S.S. (Eds.), Snakes: ecology and evolutionary biology. McGraw-Hill Publishing, New York.

Vitt L.J., Magnusson W.E., Ávila-Pires T.C.S., Lima A.P. 2008. Guia de lagartos da Reserva Adolfo Ducke, Amazônia Central. Áttema Design Editorial, Manaus.



Figura 1. *Mastigodryas boddaerti* UFOPA-H 3352 found dead after feeding on *Hemidactylus mabouia*. A - *Mastigodryas boddaerti* with *Hemidactylus mabouia* expelled from the side of the body. B - Dorsal and ventral views of *Mastigodryas boddaerti*. C - Dorsal and ventral view of *Hemidactylus mabouia*.

Table 1. Records of species predated by *Mastigodryas boddaerti*.

Táxon	Author
AMPHIBIA	
Hylidae	
<i>Scinax nebulosus</i> (Spix, 1824)	Siqueira et al. (2012)
<i>Scinax</i> sp.	Martins & Oliveira (1998)
Leptodactylidae	
<i>Leptodactylus fuscus</i> (Schneider, 1799)	Forti & Kawashita-Ribeiro (2008); Siqueira et al. (2012)
<i>Leptodactylus mystaceus</i> (Spix, 1824)	Siqueira et al. (2012)
<i>Leptodactylus longirostris</i> Boulenger, 1882	Martins & Oliveira (1998)
REPTILIA	
Reptile eggs	Siqueira et al. (2012)
Gekkonidae	
<i>Hemidactylus mabouia</i> (Moreau de Jonnès, 1818)	Present study
<i>Hemidactylus</i> sp.	Siqueira et al. (2012)
Scincidae	
<i>Copeoglossum</i> sp.	Siqueira et al. (2012)
Sphaerodactylidae	
<i>Gonatodes humeralis</i> (Guichenot, 1855)	Martins & Oliveira (1998); Siqueira et al. (2012)
<i>Gonatodes</i> sp.	Siqueira et al. (2012)
Teiidae	
<i>Ameiva ameiva</i> (Linnaeus, 1758)	Bernarde & Abe (2010); Siqueira et al. (2012); Eversole et al. (2023)
<i>Cnemidophorus lemniscatus</i> (Linnaeus, 1758)	Martins & Oliveira (1998); Siqueira et al. (2012)
<i>Cnemidophorus</i> sp.	Martins & Oliveira (1998)
<i>Kentropys calcarata</i> Spix, 1825	Martins & Oliveira (1998)
<i>Kentropyx</i> sp.	Siqueira et al. (2012)
<i>Tupinambis</i> sp.	Siqueira et al. (2012)
Tropiduridae	
<i>Uranoscodon superciliosus</i> (Linnaeus, 1758)	Siqueira et al. (2012)
<i>Tropidurus</i> sp.	Siqueira et al. (2012)

Gymnophthalmidae	
<i>Arthrosaura</i> sp.	Martins & Oliveira (1998)
AVES	
Dendrocolaptidae	
<i>Glyphorhynchus</i> sp.	Beebe (1946)
Troglodytidae	
<i>Troglodytes musculus</i> Naumann, 1823	Siqueira et al. (2012)
MAMMALIA	
Rodent	Siqueira et al. (2012)
INSECTA	
Caelifera	Beebe (1946)



Phimophis guerini
Brasília, DF
@Afonso Meneses



Boana ericae
Alto Paraíso de Goiás, GO
@Afonso Meneses