

First record of ophiophagy in *Erythrolamprus poecilogyrus* (Wied, 1825) (Serpentes: Dipsadidae)

Omar Machado Entiauspe-Neto^{1,*}, Anderson Maciel Rocha² & Daniel Loebmann¹

¹ Universidade Federal do Rio Grande, Instituto de Ciências Biológicas, Laboratório de Vertebrados. Avenida Itália, s/nº, km 8, Carreiros, CEP 96203-900, Rio Grande, Rio Grande do Sul, Brasil.

² Faculdades Cathedral de Ensino Superior, Faculdade Cathedral, Laboratório de Zoologia Aplicada de Vertebrados Terrestres e Aquáticos. Avenida Luis Canuto Chaves, 293, Caçari, CEP 69307-655, Boa Vista, RR, Brasil.

* Corresponding author email: omarentiauspe@hotmail.com.

The Neotropical dipsadid snake *Erythrolamprus poecilogyrus* (Wied, 1825) is widely distributed in the Neotropical region, occurring in most of South America east of the Andes, ranging from Argentina to the Venezuela and Guyana border (Dixon & Markezich, 1992; Natera *et al.*, 2015). This species presents a terrestrial, diurnal and nocturnal activity, preying upon anurans, lizards, and fishes (Marques *et al.*, 2001; Pinto & Fernandes, 2004; Martins *et al.*, 2008; Prieto *et al.*, 2012).

During fieldwork conducted at 3 June 2016 at 22:30 h, with 23°C air temperature, in the locality of Água Boa, Boa Vista municipality (02°49'12"N; 60°40'19"W), state of Roraima, Brazil, we observed a female individual of *Erythrolamprus poecilogyrus schotti* (Schlegel, 1837) (CHFURG 5068, Figure 1A) during a head first predation of a female *Leptodeira annulata* (Linnaeus, 1758) (CHFURG 5067, Figure 2B).

After its collection, the individuals were fixed in formalin 10% and preserved in ethanol 70%, being subjected to measurement and dissection of its stomach contents, where the *E. poecilogyrus schotti* individual presented a tail length of 110 mm, head length of 18.5 mm and body length of 460 mm, containing the *L. annulata* (Linnaeus, 1758) (CHFURG 5067, Figure 2B) individual in its stomach; the *L. annulata* individual presented three eggs in its interior, a damaged tail of approximately 110 mm, head length of 26.6 mm and body length of 420 mm.

Our observations represent the first record of ophiophagy for *E. poecilogyrus*, suggesting that it may present opportunistic predatory behavior, considering that it inhabits a wide array of biomes throughout its distribution and therefore, suitable foraging stratum, thus leading individuals to consume any putatively suitable preys. We believe that this pattern could be reflected in other species of the genus, as Bonfligio & Lema (2006) reported ophiophagy of *Erythrolamprus miliaris* Linnaeus, 1758 on *Helicops infrataeniatus* Jan, 1865, while Pereira *et al.* (2006) reported cannibalism in the same species. Further observations of natural history events are warranted in order to increase our knowledge of its feeding habits.

ACKNOWLEDGEMENTS

OME-N thanks CNPQ for the PIBIC Grant (136628/2016-8). Fieldwork was carried and specimens were collected under IBAMA permit 43658 granted to D. Loebmann.




Figure 1. A: Specimen of *Erythrolamprus poecilogyrus* (CHFURG 5068); B: Specimen of *Leptodeira annulata* (CHFURG 5067).

REFERENCES

- Bonfiglio, F. & Lema, T. 2006.** Ofiofagia em *Liophis miliaris* (Serpentes, Colubridae). *Biociências*, 14(2):221-222.
- Dixon, J. R. & Markezich, A. L. 1992.** Taxonomy and geographic variation of *Liophis poecilogyrus* (Wied) from South America (Serpentes: Colubridae). *Tex. J. Sci.*, 44(2):131-136.
- Marques, O. A. V., Eterovic, A. & Sazima, I. 2001.** *Serpentes da Mata Atlântica: guia ilustrado para a Serra do Mar*. Holos, Ribeirão Preto. 184 pp.
- Martins, M., Marques, O. A. V. and Sazima, I. 2008.** How to be arboreal and diurnal and still stay alive: microhabitat use, time of activity, and defense in Neotropical forest snakes. *South American Journal of Herpetology*, 3:58-67.
- Natera-Mumaw, M., Esqueda González, L. F. & Castelaín Fernandez, M. 2015.** *Atlas Serpientes de Venezuela. Una Visión Actual de su Diversidad*. Santiago, Chile, pp. 456.
- Pereira, H. B., Hess, P., Teixeira, M. & Domingues M. F. 2006.** *Liophis miliaris* (common water snake): Cannibalism. *Herp. Bull.*, 97:36-37.
- Pinto, R. R. & Fernandes, R. 2004.** Reproductive biology and diet of *Liophis poecilogyrus poecilogyrus* (serpentes, Colubridae) from southeastern in Brazil. *Phyllomedusa*, 3(1):9-14.
- Prieto, Y. A., Giraud, A. R. & López, M. S. 2012.** Diet and sexual dimorphism of *Liophis poecilogyrus* (Serpentes: Dipsadidae) from the wetlands regions of Northeast Argentina. *J. of Herpetol.*, 46:402-406.
- Sawaya, R. J., Marques, O. A. V. & Martins, M. R. C. 2008.** Composição e história natural das serpentes de Cerrado de Itirapina, São Paulo, sudeste do Brasil. *Biota Neotrop.*, 8(2):127-149.



Ameiva ameiva, Guarapari, Espírito Santo. Foto: Daniel Passos.