

Predation of *Ameivula nativo* (Squamata: Teiidae) by *Gampsonyx swainsonii* (Accipitriformes: Accipitridae) in Espírito Santo, Brazil

Diogo Andrade Koski¹, Gustavo Soares de Oliveira², Aline P. Valadares Koski³, Henrique C. Costa^{4,*}

¹ Instituto de Ciências Biológicas, Políticas e Sociais "Dom Vasco Fernandes Coutinho" IVAFEC-ES. Rua Castro Alves, 80, Bairro Garoto, CEP 29120-610, Vila Velha, ES, Brasil.

² Rua Chafic Murad, 106, Bento Ferreira, CEP 29050-600, Vitória, ES, Brasil.

³ Universidade Federal do Espírito Santo, Programa de Pós-Graduação em Engenharia e Desenvolvimento Sustentável, Centro Tecnológico. Avenida Fernando Ferrari, 514, Goiabeiras, CEP 29075-910, Vitória, ES, Brasil.

⁴ Universidade Federal de Minas Gerais, Programa de Pós-Graduação em Zoologia, Departamento de Zoologia. Avenida Antônio Carlos, 6.627, Pampulha, CEP 31270-901, Belo Horizonte, MG, Brasil.

* Corresponding author: ccostah@gmail.com

Ameivula nativo (Rocha, Bergallo & Peccinini-Seale, 1997) is a unisexual lizard species endemic to restinga habitats (coastal sand dunes) in Espírito Santo and Bahia states, eastern Brazilian coast (Menezes and Rocha, 2013). It was recently included in the Brazilian red list of threatened species as "Endangered", due to its habitat specificity and estimated loss of 50% of its area of occupancy, with possible extinction of some subpopulations (ICMBio, 2014; MMA, 2014). *Ameivula nativo* is also considered as "Vulnerable" in the state of Espírito Santo (Espírito Santo, 2005).

Natural history information on *Ameivula nativo* is restricted to three populations: its type-locality (Linhares, state of Espírito Santo) (Bergallo and Rocha 1993 [as *Cnemidophorus ocellifer*]); Parque Estadual Paulo César Vinha (Guarapari, Espírito Santo) (Peloso and Pavan, 2007; Peloso *et al.*, 2008); and Prado, state of Bahia (Menezes *et al.*, 2004a, b, 2008). So far, only the lizard *Tropidurus torquatus* and the snake *Philodryas patagoniensis* were recorded as predators of *A. nativo* (Peloso and Pavan, 2007). Here we record the predation of *A. nativo* by a third species, *Gampsonyx swainsonii* (Aves: Accipitriformes: Accipitridae).

The Pearl kite (*G. swainsonii*) is a monotypic kite species occurring from Nicaragua to Paraguay, Argentina and Brazil (Ferguson-Lees and Christie, 2001). Its habitats include a variety of vegetation types, from gallery forests to savannas, and natural clearings (Sick, 1997; Ferguson-Lees and Christie, 2001). The diet of Pearl kites consists of insects, small lizards, mammals and birds (Oniki, 1995; Sick, 1997; Ferguson-Lees and Christie, 2001).

The predation event was observed during an avifaunal survey at Parque Estadual Paulo César Vinha (PEPCV), a large undisturbed restinga in Guarapari, state of Espírito Santo, southeastern of Brazil (20°35' S; 40°24' W; near sea level). On 26 March 2015, 11h15, GSO was observing a Pearl kite (*G. swainsonii*) perched on a tree when suddenly the bird flew to the ground. Afterwards, the kite flew to an electric cable with a specimen of *A. nativo* in its beak (Figure 1). The lizard was eviscerated and partially eaten by the bird. About three minutes after, the kite flew away carrying the lizard in its claws.

The lizard could be identified as an individual of *A. nativo* because of its overall teiid morphology and its color pattern of olive and black with thin light stripes (Rocha *et al.*, 1997). Besides, *A. nativo* is the only species of *Ameivula* occurring in the coast of Espírito Santo (Menezes and Rocha, 2013).

Predation events concerning species of *Ameivula* (formerly part of *Cnemidophorus* [Harvey *et al.* 2012]) are largely based on *A. ocellifera*. This can be explained by *A. ocellifera* being the most widespread species of the genus and that most species of *Ameivula* were described in the last few years (Costa and Bérnills, 2014). Known predators of *Ameivula* are mostly snakes, but also birds, invertebrates, and lizards, including conspecifics (e.g., Vitt and Vangilder, 1983; Michaud and Dixon, 1989;



Figure 1: A Pearl Kite (*Gampsonyx swainsonii*), holding a lizard *Ameivula nativo* it captured in the restinga of Parque Estadual Paulo César Vinha, municipality of Guarapari, state of Espírito Santo, southeastern Brazilian coast. Photo: GSO.

Lira-da-Silva, 2009; Gogliath *et al.*, 2010; Sales *et al.*, 2010; Mesquita *et al.*, 2013; Gaiarsa *et al.*, 2013). Besides reporting for the first time a bird as predator of *A. native*, our observation adds *Gampsonyx swainsonii* as a predator of the genus *Ameivula*.

ACKNOWLEDGEMENTS

HCC is funded by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) through a doctoral fellowship.

REFERENCES

- Bergallo, H.G. and C.F. Duarte Rocha. 1993.** Activity patterns and body temperatures of two sympatric lizards (*Tropidurus torquatus* and *Cnemidophorus ocellifer*) with different foraging tactics in southeastern Brazil. *Amphibia-Reptilia*, 14:312-315.
- Costa, H.C. and R.S. Bérnils. 2014.** Répteis brasileiros: Lista de espécies. *Herpetologia Brasileira*, 3:74-84.
- Espírito Santo. 2005.** Decreto nº 1499-R, de 13 de junho de 2005. Declara as espécies da fauna e flora silvestres ameaçadas de extinção no Estado do Espírito Santo, e dá outras providências. Vitória: Diário Oficial do Estado do Espírito Santo, Poder Executivo.
- Ferguson-Lees, J. and D. A. Christie. 2001.** Raptors of the World. Houghton Mifflin, New York, 992 pp.
- Gaiarsa, M.P., L.R.V. de Alencar and M. Martins. 2013.** Natural History of Pseudoboine Snakes. *Papéis Avulsos de Zoologia*, 53:261-283.
- Gogliath, M., L.B. Ribeiro and E.M.X. Freire. 2010.** *Cnemidophorus ocellifer* (Spix's whiptail): Predation. *Herpetological Bulletin*, 114:36-38.
- Harvey, M.B., G.N. Ugueto and R.L. Gutberlet Jr. 2012.** Review of Teiid Morphology with a Revised Taxonomy and Phylogeny of the Teiidae (Lepidosauria: Squamata). *Zootaxa*, 3459:1-156.
- ICMBio – Instituto Chico Mendes de Biodiversidade. 2014.** Espécies Ameaçadas – Lista 2014. Available at: www.icmbio.gov.br/portal/biodiversidade/fauna-brasileira/lista-de-especies.html [Accessed June 08, 2015].
- Lira-da-Silva, R.M. 2009.** *Bothrops leucurus* Wagler, 1824 (Serpentes, Viperidae): Natural History Venom, and Envenomation. *Gazeta Médica da Bahia*, 79:56-65.
- Menezes, V.A. and C.F.D. Rocha. 2013.** Geographic distribution, population densities, and issues on conservation of whiptail lizards in restinga habitats along the eastern coast of Brazil. *North-Western Journal of Zoology*, 9:337-344.
- Menezes, V.A., C.F.D. Rocha and G.F. Dutra. 2004a.** Reproductive Ecology of the Parthenogenetic Whiptail Lizard *Cnemidophorus natio* in a Brazilian Restinga Habitat. *Journal of Herpetology*, 38:280-282.
- Menezes, V.A., D. Vrcibradic, J.J. Vicente, G.F. Dutra and C.F.D. Rocha. 2004b.** Helminths infecting the parthenogenetic whiptail lizard *Cnemidophorus natio* in a restinga habitat of Bahia State, Brazil. *Journal of Helminthology*, 78:323-328.
- Menezes, V.A., G.F. Dutra and C.F.D. Rocha. 2008.** Feeding habits of the endemic tropical parthenogenetic lizard *Cnemidophorus natio* (Teiidae) in a restinga area of northeastern Brazil. *Journal of Natural History*, 42:2575-2583.
- Mesquita, P.C.M.D., D.C. Passos, D.M. Borges-Nojosa and S.Z. Cechin. 2013.** Ecologia e história natural das serpentes de uma área de Caatinga no nordeste brasileiro. *Papéis Avulsos de Zoologia*, 53:99-113.
- Michaud, E.J. and J.R. Dixon. 1989.** Prey items of 20 species of the Neotropical colubrid snake genus *Liophis*. *Herpetological Review*, 20:39-41.
- MMA – Ministério do Meio Ambiente. 2014.** Portaria nº 244 de 17 de dezembro de 2014. Diário Oficial da União 245:121-126.
- Oniki, Y. 1995.** *Gampsonyx swainsonii* come uma rolinha. *Atualidades Ornitológicas*, 65:5.
- Peloso, P.L.V. and S.E. Pavan. 2007.** *Cnemidophorus natio* (NCN). Predation. *Herpetological Review*, 38:451-452.
- Peloso, P.L.V., C.F.D. Rocha, S.E. Pavan and S.L. Mendes. 2008.** Activity and microhabitat use by the endemic whiptail lizard, *Cnemidophorus natio* (Teiidae), in a restinga habitat (Setiba) in the state of Espírito Santo, Brazil. *South American Journal of Herpetology*, 3:89-95.
- Rocha, C.F.D., H.G. Bergallo and D. Peccinini-Seale. 1997.** Evidence of an Unisexual Population of the Brazilian Whiptail Lizard Genus *Cnemidophorus* (Teiidae), with Description of a New Species. *Herpetologica*, 53:374-382.
- Sales, R.F.D., L.B. Ribeiro and E.M.X. Freire. 2010.** *Cnemidophorus ocellifer* (Spix's Whiptail). Cannibalism. *Herpetological Review*, 41:217-218.
- Sick, H. 1997.** Ornitologia Brasileira. J. F. Pacheco [ed.], Editora Nova Fronteira, Rio de Janeiro, 912 pp.
- Vitt, L.J. and L.D. Vangilder. 1983.** Ecology of a Snake Community in Northeastern Brazil. *Amphibia-Reptilia*, 4:273-296.



Epictia sp., Gaúcha do Norte. Foto: Reuber Brandão.