

Predation attempt on *Dermatonotus muelleri* (Boettger, 1885) (Anura: Microhylidae) by *Caracara plancus* (Miller, 1777) (Aves: Falconidae) with a new defensive behavior reported

Daniela Sifuentes Batista^{1, 4}, Diego Gomiero Cavalheri^{2, 3*}, Sóstenes Pelegrini^{4, 5}, André Moreira Assalim⁴, Adriana Akemi Kuniy⁴, Sarah Mângia³

1 Universidade Federal de Mato Grosso, 78600-000 Barra do Garças, MG, Brazil.

2 Programa de Pós-graduação em Biologia Animal, Instituto de Biociências, Letras e Ciências Exatas, Universidade Estadual Paulista Júlio de Mesquita Filho, 15054-000 São José do Rio Preto, SP, Brazil.

3 Instituto de Biociências, Universidade Federal de Mato Grosso do Sul, 79070-900 Campo Grande, MS, Brazil.

4 JGP Consultoria e Participações Ltda. R. Américo Brasiliense 615, 04715-000 São Paulo, SP, Brazil.

5 Programa de Pós-Graduação em Conservação de Fauna, Centro de Ciências Biológicas e da Saúde, Universidade Federal de São Carlos, 13565-905 São Carlos, SP, Brazil.

*Corresponding author. E-mail: diego.cavalheri@unesp.br

DOI: [10.5281/zenodo.5211450](https://doi.org/10.5281/zenodo.5211450)

Amphibians are important components of the trophic networks in many ecosystems, in different life stages, as prey for both vertebrates and invertebrates (Toledo 2005, Toledo et al. 2007, Haddad et al. 2013). Although some animals eat frogs opportunistically (e.g. Delaix-Zaqueo et al. 2017, Cavalcante et al. 2019, Güell et al. 2019), anurans represent a significant part of the diet of many species (e.g. Menin et al. 2005, Toledo 2005, Toledo et al. 2007, Costa

et al. 2012, Jones et al. 2021). For this reason amphibians exhibit numerous strategies to avoid predators, including stiff-legged behavior, puffing up the body, and production of harmful secretions (Toledo et al. 2011).

At 10:15h on 24 November 2020, during a wildlife rescue in São Gonçalo do Gurguéia municipality (10.115°S, 45.285°W), state of Piauí, northeastern Brazil, we observed an adult Southern Caracara, *Caracara plancus* (Falco-

nidae) manipulating an object on the ground with its foot. The action was recorded using a video camera and the video is deposited in Fonoteca Neotropical Jacques Vielliard – under the voucher ZUEC-VID 952. We identified the potential prey as a Müller's termite frog, *Dermatonotus muelleri*, a burrowing species from the diagonal belt of open formations in Paraguay, Bolivia, Argentina, and Brazil (Oliveira et al. 2018). While the Caracara attacked the frog, the frog puffed up the body (Fig. 1A). After using its foot 14 times, the Caracara pecked once (Fig. 1B) and immediately shook its head (Fig. 1C). Fifteen seconds later, the bird moved away from the prey and, one minute later, flew away. After two minutes, one of us (DSB) approached and handled the frog, which remained puffed up and appeared to have an injury on its back (Fig. 1D). Slimy glandular secretion by the skin is a mechanism of defense of amphibians, usually activated by stress or injury, and the secretions vary from odoriferous to extraordinarily toxic (Toledo & Jared 1995, Gomes et al. 2007; Vitt & Caldwell 2013). After external examination, DSB released the frog in a safe location.

The Southern Caracara is a bird of prey with a generalist diet, feeding on insects, fish, mammals, birds, reptiles, and amphibians (Vargas et al. 2007, Idoeta & Roesler 2012). Although amphibians are included in its diet (Croza-

riol & Gomes 2009), they appear to be an occasional prey item (Vargas et al. 2007). This species is known to follow agricultural machinery to eat unearthed animals (see Sazima & Augusto 1991; Zamprogno & Sazima 1993, Assis & Costa 2020). The reaction of the Caracara of shaking its head immediately after pecking the frog was probably due to the secretions from the *D. muelleri*'s skin (see Cavalcante et al. 2017). The behavior was similar to the reaction of chickens after attempting to prey on *Brachycephalus ephippium* (see supplementary videos of Rebouças et al. 2019), another toxic anuran.

This is also the first report of the defensive behavior of puffing up the body in *D. muelleri*. This behavior consists of filling the lungs with air to increase the size of the frog, discouraging the predator (Toledo et al. 2011). The association of this defensive behavior and skin secretion has been documented for other amphibians (Toledo et al. 2011, Menezes & Corrêa 2020). Combining defensive behaviors results in a higher chance of escape from potential predators that are discouraged only when multiple signals are displayed (Toledo et al. 2011). While some anurans produce skin secretion at the same time as puffing up the body (e.g. *Physalaemus nattereri*, Lenzi-Mattos et al. 2005, *Odontophrynus* spp. Borteiro et al. 2018), others use this strategy only after physical contact (e.g. *Centrolene savagei*, Esco-

bar-Lasso & Rojas-Morales 2012). For *D. muelleri*, the synergistic behavior of puffing up the body and the skin secretion was effective in repelling the Crested Caracara. Despite some records of predation on *D. muelleri* (Wild 2001, Gavira et al. 2008, Stănescu et al. 2014, Caldas et al. 2017, Leal et al. 2018, Andrade et al. 2020) little is known about how its defensive strategies act against possible predators.

ACKNOWLEDGMENTS

DGC thanks Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Funding code 001.

REFERENCES

- Andrade H., Moura S., Almeida M., José E. 2020. Diet review of *Erythrolamprus poecilogyrus* (Wied-Neuwied, 1825) (Serpentes: Dipsadidae), and first record of *Dermatonotus muelleri* (Boettger, 1885) (Anura: Microhylidae) as a prey item in Sergipe State, northeastern Brazil. *Herpetology Notes* 13:1065–1068.
- Assis C., Costa H.C. 2020. *Leposternon octostegum*: New record of an endangered worm lizard species, with comments on habitat and opportunistic predators (Squamata: Amphisbaenia). *Herpetologia Brasileira* 9:109–120.
- Borteiro C., Rosset S.D., Kolenc F., Bar-rasso D.A., Lescano J.N., Baldo D. 2018. Stereotyped defensive behaviours in frogs of the genus *Odontophrynus* (Amphibia: Anura: Odontophrynidae). *Current Herpetology* 37:172–179. doi: [10.5358/hsj.37.172](https://doi.org/10.5358/hsj.37.172)
- Caldas F.L.S., Almeida B.J.M., Santos R.A. 2017. Predation of *Dermatonotus muelleri* (Anura, Microhylidae) by *Guira guira* (Cuculiformes, Cuculidae) in the coastline of the Sergipe state, northeastern Brazil. *Pesquisa e Ensino Em Ciências Exatas e Da Natureza* 1:95–98. doi: [10.29215/pecen.v1i2.447](https://doi.org/10.29215/pecen.v1i2.447)
- Cavalcante I.D., Antoniazzi M.M., Jared C., Pires O.R., Sciani J.M., Pimenta D.C. 2017. Venomics analyses of the skin secretion of *Dermatonotus muelleri*: Preliminary proteomic and metabolomic profiling. *Toxicon* 130:127–135. doi: [10.1016/j.toxicon.2017.02.028](https://doi.org/10.1016/j.toxicon.2017.02.028)
- Cavalcante, T. Simões, P. I. Mourthe, I. 2019. Predation of *Boana boans* (Anura, hylidae) by an opportunistic frugivorous primate. *Acta Amazonica* 49:307–310. doi: [10.1590/1809-4392201901430](https://doi.org/10.1590/1809-4392201901430)
- Costa H.C., Martins L.B., Carvalho T.R. 2012. *Xenodon neuwiedii* (Serpentes: Xenodontinae): Dieta. *Herpetologia Brasileira* 1:38–40.

- Crozariol M.A., Gomes F.B.R. 2009. Predação de *Leptodactylus fuscus* (Schneider, 1799) (Anura: Leptodactylidae) por *Caracara plancus* (Miller, 1777) (Aves: Falconidae), com notas sobre história natural, no Vale do Paraíba do Sul, SP. *Atualidades Ornitológicas* 149:4–5.
- Delaix-Zaqueo K., Melo-Sampaio P.R., Oliveira I.L.A., Miranda G.K.D., Calderon L.A. 2017. The opportunistic behavior of a common predator in aquatic systems in Amazonia: predation on robber-frog *Pristimantis cf. fenestratus* by trahira *Hoplias malabaricus* (Bloch, 1794). *Herpetology Notes* 10:425–427.
- Escobar-Lasso S., Rojas-Morales J. 2012. Antipredatory Behaviors of the Colombian Endemic Glassfrog *Centrolene savagei* (Anura: Centrolenidae). *Boletín Científico Museo de Historia Natural* 16:226–232.
- Gavira R.S.B., Bovo R.P., Andrade D.V. 2008. *Rhinella schneideri*. Diet. *Herpetological Review* 44:657.
- Gomes A., Giri B., Saha A., Mishra R., Dasgupta S.C., Debnath A., Gomes A. 2007. Bioactive molecules from amphibian skin: Their biological activities with reference to therapeutic potentials for possible drug development. *Indian Journal of Experimental Biology* 45:579–593.
- Güell B.A., González K., Pedroso-Santos F. 2019. Opportunistic predation by two aquatic-feeding predators on an explosive-breeding aggregation of arboreal gliding treefrogs (*Agalychnis spurrelli* Boulenger, 1913; Anura: Phyllomedusidae) on the Osa Peninsula of Costa Rica. *Herpetology Notes* 12:795–798.
- Haddad C.F.B., Toledo L.F., Prado C.P.A., Loebmann D., Gasparini J.L.L., Sazima, I. 2013. Guia dos anfíbios da Mata Atlântica: Diversidade e biologia. Editora Anolisbooks, São Paulo.
- Idoeta F.M., Roesler I. 2012. Presas consumidas por el Carancho (*Caracara plancus*) durante el periodo reproductivo, en el noroeste de la provincia de Buenos Aires. *Nuestras Aves* 57:80–83.
- Jones P.L., Divoll T.J., Dixon M.M., Aparicio D., Cohen G., Mueller U.G., ... Page R.A. 2021. Sensory ecology of the frog-eating bat, *Trachops cirrhosus*, from DNA metabarcoding and behavior. *Behavioral Ecology* 31:1420–1428. doi: [10.1093/beheco/araa100](https://doi.org/10.1093/beheco/araa100)
- Leal E.S.B., Chaves L.S., Prado-Neto J.G., Passos-Filho P.B., Ramalho D.F., Guerra-Filho D.Q., ... Moura G.J.B. 2018. What constitutes the menu of *Trachops cirrhosus* (Chiroptera)? a review of the species' diet. *Neotropical Biology and Conservation* 13:337–346. doi: [10.4013/nbc.2018.134.08](https://doi.org/10.4013/nbc.2018.134.08)

- Lenzi-Mattos R., Antoniazzi M.M., Haddad C.F.B., Tambourgi D.V., Rodrigues M.T., Jared C. 2005. The inguinal macroglands of the frog *Physalaemus nattereri* (Leptodactylidae): Structure, toxic secretion and relationship with deimatic behaviour. *Journal of Zoology* 266:385–394.
- Meneses A.S.O., Corrêa B.A.A.P. 2020. Defensive behaviors in two *Proceratophrys* species (Anura: Odontophrynidae) from central Brazilian Cerrado. *Cuadernos de Herpetología* 34:265–269. doi: [10.31017/cdh.2020.\(2019-039\)](https://doi.org/10.31017/cdh.2020.(2019-039))
- Menin M., Rodrigues D.J., Azevedo C.S. 2005. Predation on amphibians by spiders (Arachnida, Araneae) in the Neotropical region. *Phyllomedusa* 4:39–47. doi: [10.11606/issn.2316-9079.v4ip39-47](https://doi.org/10.11606/issn.2316-9079.v4ip39-47)
- Oliveira E.F., Gehara M., São-Pedro V.A., Costa G.C., Burbrink F.T., Colli G.R., Rodrigues M.T., Garda A.A. 2018. Phylogeography of Muller's termite frog suggests the vicariant role of the Central Brazilian Plateau. *Journal of Biogeography* 45:2508–2519. doi: [10.1111/jbi.13427](https://doi.org/10.1111/jbi.13427)
- Rebouças R., Carollo A.B., Freitas M.O., Lambertini C. Santos, R.M.N., Toledo L.F. 2019. Is the conspicuous dorsal coloration of the atlantic forest pumpkin toadlets aposematic? *Salamandra* 55:39–47.
- Sazima I., Augusto S.A. 1991. Habits of five brazilian snakes with coral-snake pattern, including a summary of defensive tactics. *Studies on Neotropical Fauna and Environment* 26:159–164. doi: [10.1080/01650529109360848](https://doi.org/10.1080/01650529109360848)
- Stănescu F., Marangoni F., Reinko I. 2014. Predation of *Dermatonotus muelleri* (Boettger 1885) by *Lepidobatrachus llanensis* Reig and Cei 1963. *Herpetology Notes* 7:683–684.
- Toledo L.F., Ribeiro R.S., Haddad C.F.B. 2007. Anurans as prey: An exploratory analysis and size relationships between predators and their prey. *Journal of Zoology* 271:170–177. doi: [10.1111/j.1469-7998.2006.00195.x](https://doi.org/10.1111/j.1469-7998.2006.00195.x)
- Toledo L.F., Sazima I., Haddad C.F.B. 2011. Behavioural defences of anurans: an overview. *Ethology Ecology & Evolution* 23:1–25. doi: [10.1080/03949370.2010.534321](https://doi.org/10.1080/03949370.2010.534321)
- Toledo L.F. 2005. Predation of juvenile and adult anurans by invertebrates: Current knowledge and perspectives. *Herpetological Review* 36:395–400.
- Toledo R.C., Jared C. 1995 Cutaneous granular glands and amphibian venoms. *Comparative Biochemistry and Physiology* 111:1–29.

Vargas R.J., Bó M.S., Favero M. 2007. Diet of the Southern Caracara (*Caracara plancus*) in Mar Chiquita Reserve, Southern Argentina. *Journal of Raptor Research* 41:113–121.

Vitt L.J., Caldwell J.P. 2013. Herpetology: An Introductory Biology of Amphibians and Reptiles. Academic Press. Fourth Edition. doi: [10.1016/C2010-0-67152-5](https://doi.org/10.1016/C2010-0-67152-5)

Wild E. R. 2001. *Ceratophrys cranwelli* (Cranwell's Horned) Frog. Predation. *Herpetological Review* 32:102.

Zamprogno C., Sazima I. 1993. Vertebrate predation on the neotropical Amphisbaenian *Leposternon wucherer*. *Herpetological Review* 24:82–83.

Editor: Henrique C. Costa




Figure 1. Sequence of events during a predation attempt on *Dermatoneotus muelleri* by *Caracara plancus*. A) an adult *C. plancus* manipulating the frog (indicated with a red arrow) with its foot; B) the *C. plancus* pecking the *D. muelleri*; C) *C. plancus* shaking its head and moving away from the frog; D) an injury on the dorsum of the *D. muelleri*, indicated by a red arrow.