

New record and geographic distribution of *Dipsas indica* Laurenti, 1768 in the state of Ceará, northeastern Brazil

Rodrigo Castellari Gonzalez^{1,*}, Thabata Cavalcante¹, Francisco Robson Figueiredo da-Costa¹, Daniel Cassiano-Lima^{1,2}

1 Museu de História Natural do Ceará Prof. Dias da Rocha, Universidade Estadual do Ceará. 62770-000 Pacoti, CE, Brazil.

2 Centro de Ciências da Saúde, Universidade Estadual do Ceará. 60714-903 Fortaleza, CE, Brazil.

*Corresponding author. E-mail: rodcastgon@gmail.com

DOI: [10.5281/zenodo.13308172](https://doi.org/10.5281/zenodo.13308172)

D*ipsas indica* is a small dip-sadid that reaches approximately 1000 mm in length, has aglyphous dentition, and nocturnal habits (Harvey & Embert, 2008; Harvey et al., 2009; Arteaga et al., 2018). Like its congeners, it is specialized for feeding on gastropods, which they remove from their shells with their sharp teeth (Ray et al., 2012). *Dipsas indica* is a fully arboreal species (Harrington et al., 2018), found in primary and secondary humid forests and even in disturbed areas (Harvey & Embert, 2008; Nogueira et al., 2019). It is distributed throughout South America, in Bolivia, Brazil, Colombia, Ecuador, Guyana, Peru, Suriname, and Venezuela. In Brazil, it exhibits a disjunct distribution pattern, in Amazonia and the Atlantic Forest (Nogueira et al., 2019). Here we

document the first record of *Dipsas indica* in a forest remnant in the state of Ceará, northeastern Brazil.

On February 15, 2022, an adult male of *Dipsas indica* (snout-vent-length: 413 mm, tail length: 167 mm, weight: 8.5 g, Fig. 1) was found by residents of the municipality of Guaramiranga (4°14'35.3" S, 38°56'16.4" W, 801 m a.s.l.), Ceará, Brazil. The specimen was collected under permit Sisbio #42737. A tissue sample was obtained from the liver, and the specimen was subsequently fixed in 10% formalin, and stored in 70% alcohol. The specimen and tissue sample are deposited in the Herpetological Collection of the Museu de História Natural do Ceará Prof. Dias da Rocha, under the number MHNCE-R482.

The municipality of Guaramiranga is part of the Baturité Mountains, referred to as a “Brejo de altitude” (highland wetland), an ecological island within the semi-arid Caatinga (Bétard et al., 2008; Moro et al., 2015). Due to its proximity to the coast and elevation, the region features a mountain microclimate with mild temperatures, high rainfall, high humidity, and dense humid forest above 400 m (Bétard et al., 2008; Moro et al., 2015; Silveira et al., 2020). Due to these characteristics, the region serves as a refuge for a rich relictual diversity, sheltering elements of the Amazon Rainforest and the Atlantic Rainforest, as well as a significant number of endemic species and with disjunct distribution (Borges-Nojosa, 2007; Moro et al., 2015; Borges-Nojosa et al., 2019).

This distribution pattern is not unique to this species. The Baturité Mountains are home to other snakes with a similar disjunct distribution, such as *Bothrops bilineatus* (Wied, 1821), *Lachesis rhombeata* (Wied-Neuwied, 1824) and *Sibon nebulatus* (Linnaeus, 1758) (Nascimento & Lima-Verde, 1989; Borges-Nojosa, 2007; Roberto & Loebmann, 2016; Nogueira et al., 2019; Cavalcante et al., 2022, Hamdan et al. 2024). All these snakes are found in fragmented populations across Amazonia, Atlantic Forest, and the relictual forests of the Baturité Mountains.

This record of *D. indica* in the region is another example of a species with an unexpected occurrence in a location that has been widely explored in the past (Borges-Nojosa, 2007; Roberto & Loebmann, 2016). A similar case occurred with the recent record of *B. bilineatus*, found under similar circumstances (Cavalcante et al., 2022), corroborating the importance of the Baturité mountains as a regional hotspot.

Compared to its known distribution (Fig. 2), our record of *D. indica* is 868.79 km in a straight line east of its easternmost reported location in Amazonia, in the municipality of Viseu, state of Pará, Brazil (01°12'07.2" S, 46°08'34.8" W) and 990.09 km in a straight line north of the northernmost record in the Atlantic Forest, in the municipality of Laje, state of Bahia, Brazil (13°10'55.2" S, 39°25'30.0" W). Nogueira et al. (2019) stated that the species lives at low elevations, without further information. Our record was from 801 m above sea level, which represents a significant increase to the elevation range of the species.

The reptile list of the state of Ceará currently comprises 117 species (SEMA, 2022; Barreto-Lima et al., 2023), and the only occurrence of the genus *Dipsas* was of a *Dipsas mikanii* Schlegel, 1837 in the south (Ribeiro et al., 2012; Nogueira et al., 2019). New records in the state, such as *Dipsas indica* (this

study), *Bothrops bilineatus* (Wied-Neuwied, 1821) (Cavalcante et al., 2022), and *Caiman latirostris* (Daudin, 1802) (Barreto-Lima et al., 2023) indicate the need for further studies and targeted conservation efforts focused on herpetofauna, which are crucial for the preservation of natural ecosystems, ensuring the long-term persistence of the entire ecosystems and their species.

ACKNOWLEDGEMENTS

We are grateful to Mr. Zé Carneiro and L.C. Nascimento Silva for their contribution to this study; to Paulo Passos (MNRJ) for laboratory support. RGC thanks Fundação Cearense de Apoio ao Desenvolvimento Científico e Tecnológico (process number #08339530/2023).

REFERENCES

- ANA (Agência Nacional de Águas). 2003. Catálogo de Metadados da ANA: Biomas - 1:5000000, Brasília. Acessível em: <https://metadados.snirh.gov.br/geonetwork/srv/api/records/905223d9-e516-41a-aebc-ae337954f029>. Acesso: 27 de maio de 2024.
- Arteaga A., Salazar-Valenzuela D., Mebert K., Peñafiel N., Aguiar G., Sánchez-Nivicela J.C., ... Torres-Carvajal O. 2018. Systematics of South American snail-eating snakes (Serpentes, Dipsa-
- dini), with the description of five new species from Ecuador and Peru. *ZooKeys* 766:79–147. doi: [10.3897/zookeys.766.24523](https://doi.org/10.3897/zookeys.766.24523)
- Barreto-Lima A.F., Rodrigues M. P., Costa F. R., Moura Filho A. E., Campos Y. L., ... Gonzalez R. C. 2023. First record of *Caiman latirostris* (Daudin, 1802) from the state of Ceará, Northeastern Brazil. *Herpetology Notes* 16:411–414.
- Bétard F., Peulvast J.P., Sales V.C. 2008. Caracterização morfoedológica de uma serra úmida no semi-árido do Nordeste brasileiro: o caso do Maciço de Baturité - CE. *Mercator* 6:107-126.
- Borges-Nojosa D.M.; Lima D.C., Borges-Leite M.J., Castro, D.P., Lima, A.V.P. 2019. Mata Atlântica do Ceará: herpetofauna ameaçada e estratégias de conservação. Pp. 144-161, in Abrahão C.R., Moura G.J.B., Freitas M.A., Escarlata-Tavares, F. (Eds.). Plano de Ação Nacional para a Conservação da Herpetofauna Ameaçada da Mata Atlântica Nordeste. ICMBio. Brasília.
- Borges-Nojosa D.M. 2007. Diversidade de Anfíbios e Répteis da Serra de Baturité, Ceará. Pp. 225–247, in Oliveira T.S., Araújo F.S. (Eds.), Diversidade e Conservação da Biota na Serra de Baturité. Edições UFC. Fortaleza.

- Cavalcante T., Freire-Filho R., Andrade-Oliveira J.A., Lima L.S., Cassiano-Lima D., Fernandes-Ferreira H., Gonzalez, R.C. 2022. An unexpected record of the Green Jararaca *Bothrops bilineatus* (Wied-Neuwied, 1821) in the state of Ceará, northeastern Brazil (Serpentes: Viperidae). *Herpetology Notes* 15:867–871.
- Hamdan B., Bonatto S.L., Rödder D., Seixas V.C., Santos R.M.F., Santana D.J., ... Zingali R.B. 2024. When a name changes everything: taxonomy and conservation of the Atlantic Bushmaster *Lachesis Daudin*, 1803 (Serpentes: Viperidae: Crotalinae). *Systematics and Biodiversity* 22:e2366215. doi: [10.1080/14772000.2024.2366215](https://doi.org/10.1080/14772000.2024.2366215).
- Harrington S.M., De Haan J.M., Shapiro L., Ruane S. 2018. Habits and characteristics of arboreal snakes worldwide: arboreality constraints body size but does not affect lineage diversification. *Biological Journal of the Linnean Society* 125:61–71. doi: [10.1093/biolinnean/bly097](https://doi.org/10.1093/biolinnean/bly097)
- Harvey M.B., Fuenmayor G.R., Caicedo-Portilla J.R., Rueda-Almonacid J.V. 2009. Systematics of the enigmatic dipsadine snake *Tropidodipsas perijanensis* Alemán (Serpentes: Colubridae) and review of morphological characters of Dipsadini. *Herpetological Monographs* 22:106–132. doi:[10.1655/08-025.1](https://doi.org/10.1655/08-025.1)
- Harvey M.B., Embert D. 2008. Review of Bolivian *Dipsas* (Serpentes: Colubridae), with comments on other South American species. *Herpetological Monographs* 22:54–105. doi:[10.1655/07-023.1](https://doi.org/10.1655/07-023.1)
- Moro M.F., Macedo M.B., Moura-Fé M.M., Castro, A.S., Costa R.C. 2015. Vegetação, unidades fitoecológicas e diversidade paisagística do estado do Ceará. *Rodriguésia* 66:717–743. doi:[10.1590/2175-7860201566305](https://doi.org/10.1590/2175-7860201566305)
- IBGE (Instituto Brasileiro de Geografia e Estatística). 2019. Biomas e Sistema Costeiro–Marinho do Brasil – 1:250000, Rio de Janeiro. Acessível em <https://www.ibge.gov.br/geociencias/informacoes-ambientais/estudos-ambientais/15842-biomas.html>. Acesso: 27 de maio de 2024.
- IBGE (Instituto Brasileiro de Geografia e Estatística). 2021. Limites dos países da América do Sul: EPSG: 4674 - SIRGAS 2000, Rio de Janeiro, Acessível em: <https://www.geoaplicada.com/dados/limites-paises-america-do-sul/>. Acesso: 27 de maio de 2024.
- Nascimento F.P., Lima-Verde, J.S. 1989. Ocorrência de ofídios de ambientes florestais em enclaves de matas úmidas do Ceará. (Ophidia: Colubridae). *Boletim do Museu Paraense Emílio Goeldi, Série Zoologia* 5:95–100.

Nogueira C.C., Argolo A.J.S., Arzamendia V., Azevedo J.A., Barbo F.B., Bernils R.S., ... Martins M. 2019. Atlas of Brazilian Snakes: Verified point-locality maps to mitigate the Wallacean shortfall in a megadiverse snake fauna. *South American Journal of Herpetology* 14:1–274. doi:10.2994/SAJH-D-19-00120.1

Ray J.M., Montgomery C.E., Mahan, H.K., Savitzky A.H., Lips K.R. 2012. Goo-eaters: diets of the Neotropical snakes *Dipsas* and *Sibon* in Central Panama. *Copeia* 2012:197–202. doi:10.1643/CH-10-100

Ribeiro S.C., Roberto I.J., Sales D.L., Ávila R.W., Almeida W.O. 2012. Amphibians and reptiles from the Araripe bioregion, Northeastern Brazil. *Salamandra* 48:133–146.

Roberto I.J., Loebmann, D. 2016. Composition, distribution patterns, and conservation priority areas for the herpetofauna of the state of Ceará, northeastern Brazil. *Salamandra* 52:134–152.

Secretaria do Meio Ambiente. 2022. Portaria nº 146/2022, de 27 de setembro de 2022. Dispõe sobre a Lista Vermelha dos Anfíbios e Répteis Continentais Ameaçados De Extinção do Ceará. *Diário Oficial da União* 195:44-46.

Silveira A.P., Loiola M.I.B., Gomes, V.S., Lima-Verde, L.W., Oliveira, T.S., Silva E.F., ... Bruno M.M.A. 2020. Flora of Baturité, Ceará: a wet island in the Brazilian semiarid. *Floresta e Ambiente* 27:e20180320. doi:10.1590/2179-8087.032018

Editor: Henrique C. Costa




Figura 1. Adult male *Dipsas indica* (MHNCE-R482), from Guaramiranga, Ceará, Brazil.




Figura 2. Distribution map of *D. indica*, data from Nogueira et al. (2019). Red rectangle represents the new record from Guaramiranga, Ceará, Brazil. Data from Nogueira et al. 2019.