

Predation Record of *Dipoglossus fasciatus* (Squamata: Diploglossidae) by *Salvator merrianae* (Squamata: Teiidae) in Caverna do Diabo State Park, São Paulo, Brazil

Cinthia Rachel Teixeira^{1*}, Carlos Roberto da Silva Moraes², Lucas Machado Botelho¹, Edelcio Muscat¹

¹ Projeto Dacnis, Estrada do Rio Escuro, 4754, Sertão das Cotias, 11680-000 Ubatuba, SP, Brazil.

² Quilombo Sapatu, 11960-000 Eldorado, SP, Brazil.

* Corresponding author. E-mail: cicinthiarachel@gmail.com

DOI: [10.5281/zenodo.13308223](https://doi.org/10.5281/zenodo.13308223)

The tegu lizard *Salvator merrianae* (Duméril & Bibron, 1839) is a member of Teiidae characterized by large size, between 1200 to 1500 mm total length and weighing over 4.5 kg (Fitzgerald, 1992; Diniz et al., 2021). This species is terrestrial and exhibits a generalist diet, consuming a range of items including plants, invertebrates, fish, amphibians, reptiles, birds, and small mammals (Sazima & D'angelo, 2013; Cano et al., 2015; Kasperoviczus et al., 2015; Hipolito & Sazima, 2018; Vieira et al., 2018; Diniz et al., 2021). Reports have also indicated instances of it preying on the eggs of ground nesting birds and reptiles (Sazima & D'angelo, 2013; Bovendorp et al., 2008; Mazzotti et al., 2014; Galetti et al., 2009) and regurgitating remains from seabirds (Muscat et al., 2016). Its habitat ranges from French Guiana to

Uruguay, encompassing Brazil, Bolivia, Paraguay, and Argentina. Within Brazil, it is found across various biomes and has a wide distribution within the Atlantic Forest (Dirksen & Riva, 1999; Bertoluci et al., 2009; Avila et al., 2013; Cano et al., 2015; Cassiali et al., 2016; Lima et al., 2021).

The lizard *Dipoglossus fasciatus* (Gray, 1831), of the family Diploglossidae, is distributed from the Peruvian Amazon to the Atlantic Forest in southern Brazil (Avila-Pires, 1995; Marques & Sazima, 2004). Measuring from 105 to 170 mm snout-vent length, it is a terrestrial and diurnal species (Avila-Pires, 1995), and may also have arboreal habits (Marques & Sazima, 2004).

On 19 March 2023, at 2:47 pm, we recorded an event of predation in-

volving *Salvator merianae* on *Diploglossus fasciatus* in a grassy area within the Caverna do Diabo State Park (24°38'08.95" S; 48°24'16.39" W). The Caverna do Diabo State Park is a conservation unit encompassing 40,219.66 hectares of Atlantic Forest in the Vale do Ribeira region of the state of São Paulo. It spans across the municipalities of Barra do Turvo, Cajati, Eldorado, and Iporanga. At the time of the encounter, *S. merianae* had almost completely swallowed the *Diploglossus fasciatus*. Only a portion of the tail and hind limbs were visible outside the mouth (Figure 1). We observed the predation for 30 minutes until the *D. fasciatus* was completely ingested. We identified the prey as *D. fasciatus* based on the visible coloration and structural patterns. The species exhibits short limbs and a round tail with a tapered tip. Its coloration is vibrant, with golden tones and dark transverse bands forming rings around the tail, as described by Avila-Pires (1995).

We conducted a literature search for *Diploglossus fasciatus* predation by *Salvator merianae* using the Google Scholar platform between 2 August 2023 and 1 September 2023. The search comprised the following terms: predação OR presa OR predador OR depredación OR predación OR depredador OR predation OR prey OR predator OR diet OR dieta OR *Salvator merianae* OR *Tupinambis merianae* OR

Tupinambis teguixin (former name of this species [Avila-Pires, 1995]) AND "*Diploglossus fasciatus*". Grey literature (technical reports, abstracts from events, monographs, dissertations, and theses) was not considered. Additionally, we consulted compilations of squamate predation, such as Schalk & Cove (2018). We encountered literature documenting Tegu lizards preying on lizards from the families Anguidae, Iguanidae, and Polychrotidae (Offner et al., 2021); Scincidae (Silva-Jr et al., 2005, Gaiotto et al., 2021); Teiidae (Silva et al., 2014); and Tropiduridae (Silva et al., 2013). However, there were no records of predation on diploglossids, making this a novel finding for the diet of *S. merianae*.

This may also be the first documented record of predation on the species *D. fasciatus*. In the literature, records of predation involving lizards of the genus *Diploglossus* as prey are extremely rare. Only one likely event of predation of *D. monotropis* (Kuhl, 1820) by *Eurypyga helias* (Selby, 1840) (Aves: Eurypygiidae) was described by Acosta-Chaves & Zuñiga (2013) in Costa Rica. The scarcity of these predation events may be related to the lack of information about the natural history of *Diploglossus*, due to its discreet habits (Solórzano, 2001). On the other hand, *D. fasciatus* is reported to mimic *Rhinocricus* sp., a toxic millipede (Shear, 2015), which may discourage potential predators.

Although *D. fasciatus* is best known as inhabiting forested areas, it can also be found at forest edges and urban areas (Marques & Sazima, 2004; Costa et al., 2009), and the reported predator, *Salvator merianae*, occurs in different environments (e.g. Schaumburgra et al., 2014; Muscat et al., 2016; Goetz et al., 2021). Both species are frequently observed in a forest edge area within Caverna do Diabo State Park (pers. obs.), which might have facilitated the predation event. Further field observations may unveil additional details regarding the prey-predator interaction between these species. Predation records are essential for gaining a deeper understanding of trophic relationships between species. In the case of *Diploglossus fasciatus*, data like these are crucial because it is a species with limited natural history information.

ACKNOWLEDGEMENTS

We would like to thank Elsie Rotenberg for the English review and considerations on the manuscript and Fabio Luiz Donizetti, for his contribution and assistance with the authors' communication.

REFERENCES

Acosta-Chaves V., Zuñiga J.A.S. 2013. *Diploglossus monotropis* (Costa Rican

Rainbow Striped Galliwasp). Predation. *Herpetological Review* 44:508.

Avila L.J., Martinez L.E., Morando M. 2013. Checklist of lizards and amphisbaenians of Argentina: an update. *Zootaxa* 3616:201–238. doi: [10.11646/zootaxa.3616.3.1](https://doi.org/10.11646/zootaxa.3616.3.1)

Avila-Pires T.C.S. 1995. Lizards of Brazilian Amazonia (Reptilia: Squamata). *Zoologische Verhandelingen* 299:1–706.

Bertoluci J., Canelas M.A.S., Eisemberg C.C., Palmuti C.F.S., Montingelli G.G. 2009. Herpetofauna of Estação Ambiental de Peti, an Atlantic Rainforest fragment of Minas Gerais State, southeastern Brazil. *Biota Neotropica* 9(1):147–155. doi: [10.1590/S1676-06032009000100017](https://doi.org/10.1590/S1676-06032009000100017)

Bovendorp R.S., Alvarez A.D., Galetti M. 2008. Density of the Tegu Lizard (*Tupinambis merianae*) and its role as nest predator at Anchieta Island, Brazil. *Neotropical Biology and Conservation* 3(1):9-12

Cano P.D., Ball H.A., Carpinetto M.F., Peña G.D. 2015. Reptile checklist of Río Pilcomayo National Park, Formosa, Argentina. *Check List* 11(3):1658. doi: [10.15560/11.3.1658](https://doi.org/10.15560/11.3.1658)

- Cassiali P., Scott N.J., Ortíz A.L.A., Fitzgerald L.A., Smith P. 2016. The Reptiles of Paraguay: Literature, Distribution, and an Annotated Taxonomic Checklist. *Special Publication of The Museum of Southwestern Biology* 11:1–373.
- Diniz H.S., Feio R.N., Assis C.L., Guedes J.J.M. 2021. Diet of *Salvator merianae* (Squamata: Teiidae): New prey item and review of predation records. *North-Western Journal of Zoology* 17:309–314.
- Dirksen L., Riva I. 1999. The lizards and amphisbaenians of Bolivia (Reptilia, Squamata): *Check-list, localities, and bibliography*. *Graellsia* 55:199–215.
- Galetti M., Bovendorp R.S., Fadini R.F., Gussoni C.O.A., Rodrigues M., Alvarez A.D., ... Alves K. 2009. Hyper abundant mesopredators and birds extinction in an Atlantic forest island. *Zoologia* 26:28–298. doi: [10.1590/S1984-46702009000200011](https://doi.org/10.1590/S1984-46702009000200011)
- Gaiotto J.V., Abrahão C.R., Dias R.A., Bugoni L. 2021. Diet of invasive cats, rats and tegu lizards reveals impact over threatened species in a tropical island. *Perspectives in Ecology and Conservation* 18:294–303. doi: [10.1016/j.pecon.2020.09.005](https://doi.org/10.1016/j.pecon.2020.09.005)
- Goetz S.M., Steen D.A., Miller M.A., Guyer C., Kottwitz J., Roberts J.F., ... Reed R.N. 2021. Argentine Black and White Tegu (*Salvator merianae*) can survive the winter under semi-natural conditions well beyond their current invasive range. *PLOS ONE* 16:e0245877. doi: [10.1371/journal.pone.0245877](https://doi.org/10.1371/journal.pone.0245877)
- Hipolito J.V., Sazima I. 2018. Stilts do not protect against crawlers: Limpkins preyed on by Tegu Lizards at an urban park. *Revista Brasileira de Ornitologia* 26:231–233. doi: [10.1007/BF03544436](https://doi.org/10.1007/BF03544436)
- Kasperoviczus K.N., Krüger A.C., Marques O.A. 2015. An elongated meal: the tegu lizard *Salvator merianae* eats snakes and amphisbaenians. *Herpetology Notes* 8:21–23.
- Lima J.H.A., Dias E.G., Costa R.D.L., Silva F.J., Lima E.S.M., Santos E.M., Kokubum M.N.C. 2021. Lizards and snakes of Refúgio de Vida Silvestre Matas do Siriji, an Atlantic Forest hotspot of the Pernambuco Endemism Center, Northeastern Brazil. *Biota Neotropica* 21:e20201106. doi: [10.1590/1676-0611-BN-2020-1106](https://doi.org/10.1590/1676-0611-BN-2020-1106)
- Marques O.A.V., Sazima I. 2004. História natural dos répteis da Estação Ecológica Juréia-Itatins. Pp. 257-277, in Marques O.A.V., Duleba W. (Eds.). Estação Ecológica Juréia-Itatins: ambiente físico, flora e fauna. Holos, Belo Horizonte.

- Mazzotti F.J., McEachern M., Rochford M., Reed R.N., Eckles J.K., Vinci J., Edwards J., Wasilewski J. 2014. *Tupinambis merianae* as nest predators of crocodylians and turtles in Florida, USA. *Biological Invasions* 17:47–50. doi: [10.1007/s10530-014-0730-1](https://doi.org/10.1007/s10530-014-0730-1)
- Muscat E; Olmos F; Rotenberg E.L. 2016. *Salvator merianae* (black and white tegu) scavenging around the nests of *Fregata magnificens* (frigatebird). *The Herpetological Bulletin* 135:36–37.
- Offner M.T., Campbell T.S., Johnson S.A. 2021. Diet of the invasive Argentine Black and White Tegu in central Florida. *Southeastern Naturalist* 20:319–337. doi: [10.1656/058.020.0210](https://doi.org/10.1656/058.020.0210)
- Sazima I., D'angelo G.B. 2013. Range of animal food types recorded for the tegu lizard (*Salvator merianae*) at an urban park in South-eastern Brazil. *Herpetology Notes* 6:427–430.
- Schaumburga L.G., Poletta G.L., Siroski P.A., Mudry M.D. 2014. Spontaneous genetic damage in the tegu lizard (*Tupinambis merianae*): The effect of age. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis* 15:5–9. doi: [10.1016/j.mrgentox.2014.03.007](https://doi.org/10.1016/j.mrgentox.2014.03.007)
- Shear, William A. 2015. The chemical defences of millipedes (Diplopoda): Biochemistry, physiology, and ecology. *Biochemical Systematics and Ecology* 61:78–117. doi: [10.1016/j.bse.2015.04.033](https://doi.org/10.1016/j.bse.2015.04.033)
- Silva E.A.P; Santos T.D; Leite G.N; Ribeiro L.B. 2013. *Tropidurus hispidus* (Squamata: Tropiduridae) and *Leptodactylus* cf. *fuscus* (Anura: Leptodactylidae) as prey of the teiid lizards *Salvator merianae* and *Ameiva ameiva*. *Herpetology Notes* 6:51–53.
- Silva J.S., El-Deir A.C.A., Moura G.J.B., Alves R.R.N., Albuquerque U.P. 2014. Traditional ecological knowledge about dietary and reproductive characteristics of *Tupinambis merianae* and *Hoplias malabaricus* in semiarid Northeastern Brazil. *Human Ecology* 42:901–911. doi: [10.1007/s10745-014-9698-9](https://doi.org/10.1007/s10745-014-9698-9)
- Silva-Jr J.M., Péres-Jr A.K., Sazima, I. 2005. *Euprepis atlanticus* (Noronha Skink). Predation. *Herpetological Review* 36:33–36.
- Solórzano, A. 2001. Great fire lizard. *Fauna* 2:8–12.
- Vieira R.C., Oliveira A.S., Fagundes N.J.R., Verrastro L. 2018. Predation of *Conepatus chinga* (Molina, 1782) (Carnivora, Mephitidae) by *Salvator merianae* (Duméril and Bibron, 1839)

(Squamata, Teiidae) in a field area of the state of Rio Grande do Sul, Brazil.

Herpetology Notes 11:349–351.

Editor: Henrique C. Costa



Figura 1. *Salvator merianae* preying on a *Dipoglossus fasciatus* lizard in in Caverna do Diabo State Park, São Paulo, Brazil.